

PUBLIKATIONSLISTE

Dr. Andrea Vera Keck

Stand: 22. August 2018

Novel serum markers of bone resorption: clinical assessment and comparison with established urinary indices.

Woitge HW, Pecherstorfer M, Li Y, Keck AV, Horn E, Ziegler R, Seibel MJ.
J Bone Miner Res. 1999 May;14(5):792-801.

Comparative tolerability of drug therapies for hypercalcaemia of malignancy.

Zoer N, Keck AV, Pecherstorfer M.
Drug Saf. 1999 Nov;21(5):389-406. Review.

Cancerous Bone Pain - Pathophysiology and Quantification

Pecherstorfer M, Keck A-V
Osteologie Band 2, 1999 Verlag Hans Huber AG, Bern

Effect of first treatment with aminobisphosphonates pamidronate and ibandronate on circulating lymphocyte subpopulations.

Pecherstorfer M, Jilch R, Sauty A, Horn E, Keck AV, Zimmer-Roth I, Thiebaud D.
J Bone Miner Res. 2000 Jan;15(1):147-54.

Oral Alendronate for Treatment of Charcot Neuroosteoarthropathy

(Behandlung der Charcotschen Neuroosteoarthropathie mit dem Bisphosphonat Alendronat)
B. Hammerl, J. Haller, A.V. Keck, M. Pecherstorfer
Osteologie 129-132, 2000 Verlag Hans Huber AG, Bern

Serum bone sialoprotein as a marker of tumour burden and neoplastic bone involvement and as a prognostic factor in multiple myeloma.

Woitge HW, Pecherstorfer M, Horn E, Keck AV, Diel IJ, Bayer P, Ludwig H, Ziegler R, Seibel MJ.
Br J Cancer. 2001 Feb 2;84(3):344-51.

Biochemical markers of bone formation in patients with plasma cell dyscrasias and benign osteoporosis.

Woitge HW, Horn E, Keck AV, Auler B, Seibel MJ, Pecherstorfer M.
Clin Chem. 2001 Apr;47(4):686-93.

Regional cerebral blood flow in patients suffering from post-traumatic stress disorder.
Mirzaei S, Knoll P, Keck A, Preitler B, Gutierrez E, Umek H, Köhn H, Pecherstorfer M.
Neuropsychobiology. 2001;43(4):260-4.

Beneficial effect of high-dose chemotherapy in multiple myeloma patients with unfavourable prognostic features.
Kaufmann H, Ackermann J, Greinix H, Nösslinger T, Gisslinger H, Keck A, Ludwig H, Worel N, Kalhs P, Zielinski C, Drach J.
Ann Oncol. 2003 Nov;14(11):1667-72.

Comparison of Technetium-99m-MIBI imaging with MRI for detection of spine involvement in patients with multiple myeloma.
Mirzaei S, Filipits M, Keck A, Bergmayer W, Knoll P, Koehn H, Ludwig H, Pecherstorfer M.
BMC Nucl Med. 2003 Dec 11;3(1):2.

Knochenstoffwechsel bei malignen Erkrankungen
Keck A-V, Pecherstorfer M
Journal für Mineralstoffwechsel 2003; 10 (2), 6-11

DNA-methylation analysis identifies the E-cadherin gene as a potential marker of disease progression in patients with monoclonal gammopathies.
Seidl S, Ackermann J, Kaufmann H, Keck A, Nösslinger T, Zielinski CC, Drach J, Zöchbauer-Müller S.
Cancer. 2004 Jun 15;100(12):2598-606.

Cytogenetic patterns in multiple myeloma after a phase of preceding MGUS.
Kaufmann H, Ackermann J, Odelga V, Sagaster V, Nösslinger T, Pfeilstöcker M, Keck A, Ludwig H, Gisslinger H, Drach J.
Eur J Clin Invest. 2008 Jan;38(1):53-60.

Local therapy of stage IV renal cell cancer: a case report.
Ferner M, Keck AV, Niedersüß-Beke D, Strasser-Weippl K.
Memo. accepted: Nov 2017

EGFR mutations in cell-free plasma DNA from patients with advanced lung adenocarcinomas: improved detection by Droplet Digital PCR.
Buder A, Setinek U, Hochmair MJ, Schwab S, Kirchbacher K, Keck A, Burghuber OC, Pirker R, Filipits M,
Lung Cancer. submitted: Apr 2018